



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the study and toil of ages. In all this he holds the noblest of all possessions, a brave, intelligent, and trusting wife, whose sympathy and encouragement is a constant incentive to him to work on, to penetrate still deeper in the hidden mysteries of his laborious science. He is tall, being five feet eleven inches in height, and weighs 185 pounds. He has light grayish blue eyes and a face which leaves the impression of power and capacity.

[Erratum. Page 186, beginning of 16th line from the top of page, insert "to accept the presidency".]

SOME NOVEL AND INTERESTING FORMULAS.

By J. W. NICHOLSON, A. M., LL. D., Member of the London, and New York Mathematical Societies,
and President and Professor of Mathematics, Louisiana State University, Baton Rouge, Louisiana.

These formulas are given without demonstration, thinking that their deduction would occupy more space than they probably deserve.

$$(2a+2b)^2 + a^2 + b^2 = (2a+b)^2 + (a+2b)^2 \dots\dots(1).$$

This is a simple formula for finding three square numbers whose sum is equal to the sum of two squares. Thus, for $a=5$, $b=3$, we have

$$16^2 + 5^2 + 3^2 = 13^2 + 11^2 \dots\dots(2).$$

$$(3a+3b)^n + (2a+4b)^n + a^n + b^n = (3a+4b)^n + (a+3b)^n + (2a+b)^n \dots\dots(3),$$

where $n=3, 2$ or 1 .

Thus, for $a=5$, $b=3$, we have $24^n + 22^n + 5^n + 3^n = 27^n + 14^n + 13^n \dots\dots(4)$,
where $n=3, 2$ or 1 .

$$(5a+10b)^n + (4a+11b)^n + (3a+5b)^n + (2a+8b)^n + (3a+3b)^n + (2a+6b)^n + a^n + b^n = (5a+11b)^n + (4a+6b)^n + (3a+10b)^n + (3a+8b)^n + (a+5b)^n + (2a+3b)^n + (2a+b)^n \dots\dots(5),$$

where $n=5, 4, 3, 2$ or 1 .

Thus for $a=5$, $b=2$, we have

$$45^n + 42^n + 26^n + 25^n + 22^n + 21^n + 5^n + 2^n = 47^n + 35^n + 32^n + 31^n + 16^n + 15^n + 12^n \dots\dots(6),$$

where $n=5, 4, 3, 2$ or 1 .

In (5) for $a=8$, $b=3$, we find $15^n + 10^n + 9^n + 6^n = 14^n + 13^n + 7^n + 3^n + 2^n + 1^n \dots\dots(7)$,
where $n=5, 3$ or 1 .

$$(a+32)^n + (a+24)^n + (a+18)^n + (a+10)^n + (a+4)^n + (a-4)^n + (a-10)^n + (a-18)^n + (a-24)^n + (a-32)^n = (a+30)^n + (a+28)^n + (a+16)^n + (a+8)^n + (a+6)^n + (a-6)^n + (a-8)^n + (a-16)^n + (a-28)^n + (a-30)^n \dots\dots(8),$$

where $n=5, 4, 3, 2$ or 1 .

In (8) by making $a=7$, we find

$$39^n + 31^n + 21^n + 9^n = 37^n + 35^n + 15^n + 13^n \dots\dots(9), \quad \text{where } n=5, 3 \text{ or } 1.$$

$$j = \lfloor \frac{n}{2} \rfloor = n^n - n(n-1)^n + \frac{n(n-1)}{2}(n-2)^n - \frac{n(n-1)(n-2)}{2 \cdot 3}(n-3)^n + \&c. \dots\dots(10),$$

where n is any positive integer.

Thus, for $n=5$, we have

$$1.2.3.4.5=5^5-5(4)^5+10(3)^5-10(2)^5+5(1)^5\ldots(11).$$

Some new and interesting properties of prime numbers.

If $n+1$ is a prime number, then

$$(a+nb)^x+[a+(n-1)b]^x+[a+(n-2)b]^x+\ldots(a+b)^x+a^x=m(n+1)\ldots(12),$$

where m is an integer and x any integer less than n .

In (12) for $a=0$ and $b=1$, we have

$$n^x+(n-1)^x+(n-2)^x\ldots 2^x+1=m(n+1)\ldots(13).$$

Thus, 11 will exactly divide

$$10^x+9^x+8^x+7^x+6^x+5^x+4^x+3^x+2^x+1\ldots(14) \text{ where } x=9, 8, 7, 6, 5, 4, 3, 2, 1.$$

The converse of formulas (12) or (13) is not always true, but the following are true only when $n+1$ is a prime number.

$$(a+n)^n+(a+n-1)^n+(a+n-2)^n+\ldots a^n+1=m(n+1)\ldots(15).$$

Making $a=0$, we have

$$n^n+(n-1)^n+(n-2)^n\ldots 1^n+1=m(n+1)\ldots(16).$$

That is, S_n+1 is divisible by $n+1$ when it is a prime number and only when it is prime. So far as I know this furnishes an entirely *new criterion of prime numbers*.

NOTE. The preceding formulas are taken from a paper, by the author, on "The n th power of any number expressed as the sum of the n th powers of other numbers, n being any positive integer;" which was read before the New York Mathematical Society, Dec. 3d, 1892.

NON-EUCLIDEAN GEOMETRY: HISTORICAL AND EXPOSITORY.

BY GEORGE BRUCE HALSTED, A. M., (Princeton) Ph. D., (Johns Hopkins); Member of the London Mathematical Society; and Professor of Mathematics in the University of Texas, Austin, Texas.

CHAPTER SECOND.

THE FIRST TREATISE ON NON-EUCLIDEAN GEOMETRY.

[Continued from the May Number.]

PROPOSITION I. *If two equal straight lines [sects] (fig 1.) AC , BD , make with the straight AB angles equal toward the same parts: I say that the angles at the join CD will be mutually equal.*

PROOF. Join AD , CB . Then consider the triangles CAB , DBA . It follows (Eu. I. 4.) that the bases CB , AD will be equal.

Then consider the triangles ACD , BDC . It

